

U.S. DEPARTMENT OF COMMERCE

LUTHER H. HODGES, *Secretary*

WEATHER BUREAU

ROBERT M. WHITE, *Chief*

TECHNICAL PAPER NO. 50

**Frequency of Maximum Water Equivalent  
of March Snow Cover  
in North Central United States**

Prepared by  
Cooperative Studies Section, Office of Hydrology  
U.S. Weather Bureau  
for  
Engineering Division, Soil Conservation Service  
U.S. Department of Agriculture



WASHINGTON, D.C.

1964



## CONTENTS

	Page
1. INTRODUCTION.....	1
2. BASIC DATA.....	1
3. MAXIMUM OBSERVED VALUES.....	3
4. ESTIMATION OF MAXIMUM WATER EQUIVALENT.....	8
5. FREQUENCY ANALYSIS.....	10
6. CONSTRUCTION OF WATER-EQUIVALENT MAPS.....	10
7. DISCUSSION OF WATER-EQUIVALENT MAPS.....	12
REFERENCES.....	12

### LIST OF ILLUSTRATIONS

Figure No.	Page
1. Stations providing snow data used in this study.....	2
2. Maximum observed water-equivalent values (in.) of record for March 1-15.....	4
3. Maximum observed water-equivalent values (in.) of record for March 16-31.....	5
4. Maximum observed snow depth (in.) for March 1-15.....	6
5. Maximum observed snow depth (in.) for March 16-31.....	7
6. Curve for adjusting estimates of water equivalent obtained from equation (2).....	9
7. Comparison of adjusted estimates of water equivalent with observed values.....	9
8. Probability-interpolation diagram.....	11
9. Points for which water-equivalent values were interpolated.....	11
10. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in two years.....	13
11. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in two years.....	14
12. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in five years.....	15
13. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in five years.....	16
14. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in ten years.....	17
15. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in ten years.....	18
16. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in 25 years.....	19
17. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 25 years.....	20
18. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in 50 years.....	21
19. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 50 years.....	22
20. Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in 100 years.....	23
21. Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 100 years.....	24



# FREQUENCY OF MAXIMUM WATER EQUIVALENT OF MARCH SNOW COVER IN NORTH CENTRAL UNITED STATES

## 1. INTRODUCTION

*Authority.* This report was prepared for the Soil Conservation Service to provide generalized information for planning and design purposes in connection with its Watershed Protection and Flood Prevention Program (authorization: P.L. 566, 83d Congress, and as amended).

*Scope.* Maximum water-equivalent values of snow on the ground for the first and second halves of March are presented for probabilities of 50, 20, 10, 4, 2, and 1 percent. The region covered is north of 40° N. and between 80° and 105° W.

*Accuracy of results.* The accuracy of the estimates presented is dependent on the number of stations, quality of observations, and length of record. Water-equivalent measurements are among the most inaccurate of meteorological observations. Besides the usual observational errors, the measurements may be unrepresentative when obtained from drifted snow or from unduly exposed or sheltered sites. Also, inconsistencies in day-to-day measurements often arise from the necessity to shift measurements from one observation plot to another to obtain readings in undisturbed snow cover.

The longest records of daily water-equivalent observations for the relatively small number of stations in the study area were only 11 years, and many records were for a few years only. However, measurements of snow depth have been made at many stations for many years. Equations

based on observed water-equivalent values, associated snow depths, and other pertinent parameters were derived to estimate water equivalent for those stations and periods with snow-depth measurements only. The resulting synthetic records of water-equivalent values were then subjected to frequency analysis. The results of such analyses are not so reliable as those based on long records of water-equivalent observations if these were available. Consequently, it is reasonable to expect that more reliable results could be obtained from 10 or more additional years of such observations, especially if, in the meantime, the quality of the observations were improved and the number of observing stations increased.

*Acknowledgments.* The project was under the general supervision of J. L. H. Paulhus, Chief of the Cooperative Studies Section of the Office of Hydrology, W. E. Hiatt, Acting Director. J. F. Miller was project leader. A. H. Jennings made the preliminary investigations relating the water equivalent of snow on the ground to meteorological parameters. L. L. Weiss and L. O. Feese performed the statistical investigations. N. S. Foat supervised the collection and processing of the basic data. Coordination with the Soil Conservation Service was maintained through H. O. Ogrosky, Chief, Hydrology Branch, Engineering Division.

## 2. BASIC DATA

*Primary water-equivalent network.* Basic data for the study was obtained from the records of 61 Weather Bureau first-order stations. The locations of 52 of these stations are shown in figure 1, the other 9 stations being just outside the problem

area. Observations of water equivalent of the snow on the ground at many of these stations began in 1953, but some stations did not start until much later. A few did not start such observations until 1962. The average length of record of




FIGURE 1.—Stations providing snow data used in this study.

water-equivalent observations was 9 years. Observations of snow depth were made at many of these stations for many years before the inauguration of water-equivalent measurements. The average period of record used in this study was 44 years, which includes the period during which water-equivalent measurements were also made. Table 1 provides a breakdown of the 61 stations in the water-equivalent network by length of record of snow-depth and water-equivalent measurements used.

TABLE 1.—Stations measuring both snow depth and water equivalent grouped by length of record

Years of record used	Number of stations with data on:	
	Snow depth	Water equivalent
1-5	2	12
6-10	1	5
11-15	2	44
16-20	3	0
21-25	10	0
26-30	0	0
31-35	1	0
36-40	2	0
41-45	2	0
46-50	3	0
51-55	35	0
Totals.....	61	61

*Secondary snow-depth network.* Supplementing the water-equivalent network was another network of 463 stations (fig. 1) where snow on ground was measured daily for depth only. Data for these stations were obtained from the Weather Bureau's *Climatological Data* for the period of record subsequent to 1949. Data were tabulated for all available stations, but stations with less than 5 years of record were not used. Copies of the original records for about one-fourth of these stations were obtained for the period 1939-49. These stations were selected to provide a reasonably uni-

TABLE 2.—Stations measuring snow depth only grouped by length of record

Years of record used	Number of stations
1-5	0
6-10	123
11-15	216
16-20	23
21-25	96
26-30	0
31-35	1
36-40	0
41-45	1
46-50	0
51-55	3
Total.....	463

form geographic sampling. Grouping of stations by length of record is shown in table 2.

*Supplementary snowfall data.* In addition to the network data, use was made of the "Supplementary Snowfall Data" published in *Climatological Data*. These data were obtained from field snow surveys conducted by Weather Bureau and Corps of Engineers personnel or by station observers acting under detailed instructions. Measurements of snow depth and water equivalent at designated stations were made on Tuesdays and Fridays.

Other supplementary data were obtained from unpublished listings of snow-survey data maintained at some Weather Bureau offices for some special networks.

*Quality.* Accurate measurements of representative water-equivalent values of snow on ground are very difficult to make. The depth of snow cover may vary a great deal within short distances because of drifting, and, in some cases, the determination of a representative depth and water equivalent is practically impossible.

Measurements of snow depth and water equivalent should be made in undisturbed snow and from approximately the same site. Since measurements are made daily, these limitations require that a fairly large plot of land be available for the observations. Most of the stations measuring water

TABLE 3.—*Examples of inconsistencies in water-equivalent data*

Station..... Month, Year..... Date.....	Minneapolis, Minn. March 1962				Sioux Falls, S. Dak. Feb.-Mar. 1962			
	7	8	9	10	26	27	28	1
Max. Temp. (°F.).....	33	33	36	36	5	-2	0	9
Min. Temp. (°F.).....	20	28	21	21	-12	-18	-31	-16
Precipitation (in.).....	T	T	0	.04	.02	.01	0	0
Snowfall (in.).....	0	T	0	0.2	0.6	0.3	0	0
Snow depth (in.).....	23	22	19	17	26	26	18	18
Water equiv. (in.).....	3.2	3.2	4.7	4.7	2.4	2.4	3.3	3.3
Remarks.....	Jump in WE on 9th unexplainable.				Drop in snow depth and jump in WE on 28th unexplainable.			

equivalent are located in cities or at airports where large observational plots are rarely available. The observer is then required to shift his measurements from one small plot to another when it is no longer possible to obtain readings in undisturbed snow. Such changes in observational sites may result in appreciable differences in measurements. Measurements of snow depth and water equivalent are also subject to observational errors; e.g., erroneous readings.

Table 3 shows two examples of typical discrepancies found in snow data. Such inconsistencies are not uncommon. Those shown probably resulted from changes in observation site, but discrepancies apparently resulting from observational or typographical errors were found.

### 3. MAXIMUM OBSERVED VALUES

Maximum values of record are always of interest in a frequency study of extremes. Figures 2 and 3 show the maximum water-equivalent values of record for March 1-15, and March 16-31, respectively. The associated snow depths and years of record are also shown. The maximum observed values of water equivalent in both the first and latter halves of March varied from slightly less than 1 in. along the southern and western edge of the region to over 6 in. over northern Michigan. About two-thirds of the stations observed their maximum values in the first half of March.

Figures 4 and 5 show the maximum snow depths of record at selected stations for the two halves of March. Also shown are the associated water-equivalent values and years of record. Though these stations were selected to show the maximum observed depths in their vicinities, surrounding stations had amounts of nearly equivalent magnitude. Most of the water-equivalent values are

estimated since most of the stations did not measure water equivalent. However, values for stations making such measurements are also estimated unless the maximum observed values of snow depth occurred within that part of the record when water-equivalent measurements were made. The letter "E" identifies all estimated values. (See Sect. 4.)

Maximum observed snow depths in the first half of March (fig. 4) varied from near 1 ft. in southern Ohio to about 4 ft. in northern Michigan. Along the western edge of the region snow depths were near 18 in., except in the Black Hills, where values near 4 ft. were observed.

In the latter half of the month (fig. 5) maximum snow depths in southern Ohio and Indiana were only about 6 in., while in northern Michigan they were still about 4 ft. Values along the western edge of the region were about 1 ft., except in the Black Hills, where they were nearly 4 ft.




FIGURE 2.—Maximum observed water-equivalent values (in.) of record for March 1-15.






FIGURE 3.—Maximum observed water-equivalent values (in.) of record for March 16-31





FIGURE 4.—Maximum observed snow depths (in.) for March 1-15.



**FIGURE 5.—Maximum observed snow depths (in.) for March 16-31.**

#### 4. ESTIMATION OF MAXIMUM WATER EQUIVALENT

Since the number of stations making water-equivalent measurements was so small and their period of record so short, it was decided to supplement the observed water-equivalent data with values estimated from accumulated snow depth and other pertinent parameters. These parameters had to be limited to those that could be obtained from the meteorological observations made by the stations measuring the depth only of snow. The published daily data for most of these stations include only temperature (maximum and minimum), amount of precipitation, snow fall, and snow depth.

A comparison made between maximum observed water equivalent and water equivalent associated with maximum snow depth in the same half-month period yielded the results given in table 4. Of the 240 paired items used, 68 percent showed no difference, and 83 percent showed differences of less than 0.2 in. On the basis of this comparison, it was assumed that estimates of water equivalent for maximum snow depths would, in general, reasonably represent maximum water-equivalent values.

Various parameters and different types of relationships were tested, and two equations were eventually selected—one for snow cover less than 10 days old and one for snow cover of longer duration. Strangely enough, temperature, in degree-days, did not appear to be related significantly to maximum water equivalent. Various base values and durations were tested without success.

TABLE 4.—Distribution (number of cases) of differences between maximum observed water equivalent and water equivalent observed with maximum snow depth

Difference* (in.)	Maximum observed water equivalent (in.)						
	0-0.9	1.0-1.9	2.0-2.9	3.0-3.9	4.0-4.9	5.0-5.9	6.0-6.9
0.....	35	34	33	30	11	15	6
0.1.....	6	7	11	8	2	2	0
0.2.....	1	3	1	3	1	0	0
0.3.....		3	1	2	1	1	0
0.4.....		2	2	1	0	0	1
0.5.....		1	2	2	2	0	
0.6-1.0.....		1	1	1	3	1	
1.1-1.5.....			1	1	0		
1.6-2.0.....					0		
2.1-2.5.....					0		
2.6-3.0.....					1		

\*Maximum observed WE minus WE observed with maximum snow depth—all differences positive.

Separate relations were originally developed for both new and old snow cover in the first and second halves of March, i.e., four relationships, but the only differences noted were not appreciable, and it was decided to use one relation only for new snow cover and one for snow cover more than 10 days old.

The equation for estimating maximum water equivalent ( $WE_{max}$ ) of a snow cover less than 10 days old merely equated  $WE_{max}$  to the total precipitation ( $P$ ) falling during the period of snow accumulation leading to the maximum snow depth, or

$$WE_{max} = \Sigma P \quad (1)$$

The relation between  $WE$  observed at time of maximum snow depth and  $WE$  estimated by this formula was found to have a correlation coefficient of 0.90 and a standard error of estimate of 0.1 in. The mean value of the 78 cases of observed  $WE$  was 0.8 in.

The equation for estimating maximum  $WE$  for a snow cover more than 10 days old was

$$WE_{max} = -0.061 + 0.172(SOG_{max}) + 0.675(\Sigma P_{10}) - 0.108(SOG_{max} - SOG_{-10}) \quad (2)$$

where  $SOG_{max}$  is maximum depth of snow on ground (in.) in the first or second half of March,  $\Sigma P_{10}$  is the total precipitation (in.) for the 10 days prior to date of maximum snow depth, and  $SOG_{-10}$  is the snow depth 10 days prior to date of maximum depth ( $SOG_{max}$ ). For example, a maximum snow depth of 20 in. for the second half of March, say the 19th, and a snow depth of 25 in. on the 9th would yield  $-5$  in. for ( $SOG_{max} - SOG_{-10}$ ).

The relation between  $WE$  observed at time of maximum snow depth and  $WE$  estimated by equation (2) showed a correlation coefficient of .86 and standard error of estimate of 0.8 in. The mean of the 240 observed values was 2.5 in.

Precipitation accumulations and differences in snow depth for various time intervals, e.g., 5, 15, 30 and more days, were tried, but the 10-day interval appeared to yield the best results. More




FIGURE 6.—Curve for adjusting estimates of water equivalent obtained from equation (2).

complicated equations based on the same and other parameters, such as temperature and snowfall, were tested also but no appreciable improvements were noted. However, comparisons of observed  $WE$  and  $WE$  estimated by equation (2) indicated a tendency for the equation to yield  $WE$  values that were too high for small values and too low for high values. A correction curve (fig. 6) was therefore constructed. A comparison (fig. 7) based on 112 independent paired items of observed  $WE$  and  $WE$  estimated by means of equation (2) and adjusted by means of figure 6 indicated a correlation coefficient of .85, a standard error of estimate of 0.9 in. The mean of the 112 observed items was 3.4 in.

The parameters used in equations (1) and (2) do have good physical bases. Equation (1) for example, which would be used when maximum  $WE$  is realized from one or two recent storms rather than from an accumulation of old snow, utilizes precipitation only. This is reasonable since there is relatively little time for "aging" of the snow or loss of the precipitation.




FIGURE 7.—Comparison of adjusted estimates of water equivalent with observed values.

The parameters of equation (2) appear sound also. One would naturally expect a good relationship between maximum  $WE$  and maximum depth of snow on ground. The amount of precipitation falling in the last 10 days ( $\Sigma P_{10}$ ) is highly related to  $WE_{max}$ , perhaps because in March snowfalls are likely to have a fairly high density and contribute a great deal to  $WE_{max}$ . The difference in depth between the maximum snow on ground and that 10 days before is also significant. If the depth has decreased appreciably in the last 10 days, it means that some melting has taken place, and the snow cover is soggy and, for a given snow depth, has a greater water equivalent than when the difference is positive, which would be the case for new snow in the last 10 days.

A very important point to be kept in mind is that the interest in this study was in maximum  $WE$  values in each half of March. The equations may yield unsatisfactory results for other than maximum or near-maximum  $WE$  and for regions outside the study areas, especially where orographic or maritime influences are involved.

## 5. FREQUENCY ANALYSIS

The observed and/or estimated values of maximum  $WE$  for each station and for each half of March were subjected to various tests in order to determine the most appropriate frequency distribution. This was important because the obtaining of  $WE$  values for the smaller probabilities of exceedance represents, in effect, an extrapolation of the observed or synthesized record. The usual procedure is to seek a distribution function that satisfactorily describes the observed frequencies. The distribution function is then used to provide extrapolation to the smaller probabilities.

Several distribution functions were tested, and the choice finally narrowed down to two—the gamma and the lognormal. Comparison of plotted data with computed distribution curves indicated that the lognormal distribution was somewhat more representative than the gamma. This was

the distribution used although a  $t$ -test [1] applied to the means of the 2-percent probability levels for the two distributions indicated no differences significant at the 5-percent probability level.

The lognormal analysis was programmed for the computer to yield  $WE$  values for probabilities of 50, 20, 10, 4, 2, and 1 percent. Examination of the results for long-record stations indicated reasonable values for the range of probabilities used. In order to determine if the short-record data from recent years introduced any possible bias, data from stations with 35 years or more of record were examined. Data for the last 10 and 25 years of record from each station were compared at the 50- and 2-percent probability levels, and no significant bias was indicated. Results for short-record stations naturally tended to be somewhat erratic for the lower probabilities.

## 6. CONSTRUCTION OF $WE$ MAPS

Two 50-percent probability maps were first constructed, one for each half of March (figs. 10 and 11). These maps, based on observed and estimated  $WE$  data for all stations (tables 1 and 2), show the maximum  $WE$  of snow on ground expected to be equaled or exceeded on an average of once in two years.

The next step was the construction of the 1-percent probability maps. In order to avoid inconsistencies arising from the often erratic 1-percent  $WE$  values yielded by short-record stations, ratio maps of 1-percent  $WE$  to 50-percent  $WE$  based on long-record stations were first constructed. These ratio maps, one for each half of March, were then applied to the corresponding 50-percent  $WE$  maps to obtain the 1-percent  $WE$  maps (figs. 20 and 21).

In order to obtain a consistent interpolation between the 50-percent and 1-percent maps, the probability-interpolation diagram of figure 8 was

used for estimating the  $WE$  values for the intermediate probabilities, i.e., 20, 10, 4, and 2 percent. This diagram was derived empirically from the data for long-record stations only (35 years or more). Separate interpolation diagrams for the first and second halves of March were first constructed, but since there were no appreciable differences, the diagrams were combined.

In constructing the intermediate maps, the 50- and 1-percent  $WE$  values were first read for each grid point shown in figure 9. These two values for each grid point were then plotted on the corresponding verticals of the interpolation diagram (fig. 8), a straightedge was laid along the two points, and  $WE$  values for intermediate probabilities were read at the intersection of the straightedge and corresponding verticals. The  $WE$  values thus obtained were then plotted on the corresponding grid points on the appropriate maps, and isolines were then drawn.

the specific storm of Lake Michigan. It is  
 the value obtained for the year or upward  
 which tend to be about the same as those on the  
 general map of Lake Michigan for any given day.  
 Each such value is indicated for the  
 storm of Lake Michigan. This is possible  
 a result of the varying sizes of the lakes which  
 the high value in eastern watersheds, which  
 have the largest watershed of Lake Michigan  
 are to be taken into consideration of the average  
 of the values was placed between the  
 year and the point of area around the  
 the 100 and the 200-foot water-equivalent  
 (Figure 8) for the year of 1880. The  
 year. The year 1880 is the year in which  
 higher than would be indicated by the 100-foot  
 water-equivalent or 200-foot. However, when the  
 general map and contour lines are drawn  
 together of each of the years of the  
 and the average to be higher than the  
 the general map of Lake Michigan.

FIGURE 8.—Probability-interpolation diagram. →





FIGURE 9.—Points for which water-equivalent values were interpolated.

## 7. DISCUSSION OF *WE* MAPS

The maps of figures 10-21 show the maximum *WE* of snow on ground in the first and second halves of March for 50-, 20-, 10-, 4-, 2-, and 1-percent probabilities. In other words, the 1-percent map of figure 20 shows the maximum *WE* expected to be equaled or exceeded in the first half of March on an average of once in 100 years. The 2-percent map (fig. 18) shows the values expected to be equaled or exceeded on an average of once in 50 years; the 4-percent, 25 years; the 10-percent, 10 years; the 20-percent, 5 years; and the 50-percent, 2 years.

In general, the maps show higher *WE* values for the first half of March than for the second half. The reason may well be that appreciable melting and runoff usually take place over most of the study area during March. This depletion was not evident in the colder northern Great Lakes Region, which shows higher values for the second half of March.

One very noticeable feature of the maps is the difference between the water-equivalent values on

the opposite shores of Lake Michigan. Whereas the values indicated for the western, or windward, shore tend to be about the same as those on the western shore of Lake Huron for any given latitude, much higher values are indicated for the eastern shore of Lake Michigan. This is probably a result of the warming effect of the lake in winter. The high values in extreme northwestern Michigan to the lee of the western end of Lake Superior may be another indication of this warming effect.

Good agreement was noted between the *WE* maps and the pattern of mean annual total snowfall [2] and also the snow-loads maps presented by Thom [3] for return periods of 2, 10, 25, and 50 years. The snow loads, in general, tend to be higher than would be indicated by the *WE* for corresponding probabilities. However, since the snow-loads study considered heaviest snow loads regardless of month of occurrence, the results could be expected to be higher than the *WE* values presented herein, which are based on March values only.

## REFERENCES

1. C. E. P. Brooks and H. Carruthers, *Handbook of Statistical Methods in Meteorology*, Great Britain, Meteorological Office (M.O. 538), London, 1953, 413 pp. (pp. 64-67).
2. U.S. Weather Bureau, "Mean Annual Total Snowfall," sheet of *National Atlas of the United States*, 1960.
3. H. C. S. Thom, "Building Design for Wind and Snow Loads," *Agricultural Engineering*, vol. 44, No. 8, Aug. 1963, pp. 426-431.






FIGURE 10.—Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in two years.




FIGURE 11.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in two years.




FIGURE 12.—Maximum March 1–15 water equivalent (in.) expected to be equaled or exceeded once in five years.




FIGURE 13.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in five years.




FIGURE 14.—Maximum March 1–15 water equivalent (in.) expected to be equaled or exceeded once in ten years.




FIGURE 15.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in ten years.




FIGURE 16.—Maximum March 1–15 water equivalent (in.) expected to be equaled or exceeded once in 25 years.




FIGURE 17.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 25 years.






FIGURE 18.—Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in 50 years.




FIGURE 19.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 50 years.




FIGURE 20.—Maximum March 1-15 water equivalent (in.) expected to be equaled or exceeded once in 100 years.




FIGURE 21.—Maximum March 16-31 water equivalent (in.) expected to be equaled or exceeded once in 100 years.