

**NOAA Technical Memorandum  
NWS ER-105**



---

**DENSE FOG CLIMATOLOGY FOR THE BLUE RIDGE  
FOOTHILLS AND PIEDMONT AREAS OF THE BLACKSBURG, VA  
COUNTY WARNING AREA FOR THE PERIOD 1973 -2008**

**JAN JACKSON, KEN KOSTURA AND WILLIAM PERRY**

National Weather Service Office  
BLACKSBURG, VA

Scientific Services Division  
Eastern Region Headquarters  
Bohemia, New York  
February 2011

**NOAA TECHNICAL MEMORANDA**  
National Weather Service, Eastern Region Subseries

The National Weather Service Eastern Region (ER) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready for formal publications. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to ER personnel, and usually will not be widely distributed.

Papers 1 to 22 are in the former series, ESSA Technical Memoranda, Eastern Region Technical Memoranda (ERTM); papers 23 to 37 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 38, the papers are now part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 22 are available from the National Weather Service Eastern Region, Scientific Services Division, 630 Johnson Avenue, Bohemia, NY, 11716. Beginning with 23, the papers are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, VA 22161. Prices vary for paper copy and for microfiche. Order by accession number shown in parentheses at end of each entry.

**ESSA Technical Memoranda**

ERTM	1	Local Uses of Vorticity Prognoses in Weather Prediction. Carlos R. Dunn. April 1965.
ERTM	2	Application of the Barotropic Vorticity Prognostic Field to the Surface Forecast Problem. Silvio G. Simplicio. July 1965.
ERTM	3	A Technique for Deriving an Objective Precipitation Forecast Scheme for Columbus, Ohio. Robert Kuessner. September 1965.
ERTM	4	Stepwise Procedures for Developing Objective Aids for Forecasting the Probability of Precipitation. Carlos R. Dunn. November 1965.
ERTM	5	A Comparative Verification of 300 mb. Winds and Temperatures Based on NMC Computer Products Before and After Manual Processing. Silvio G. Simplicio. March 1966.
ERTM	6	Evaluation of OFDEV Technical Note No. 17. Richard M. DeAngelis. March 1966.
ERTM	7	Verification of Probability of Forecasts at Hartford, Connecticut, for the Period 1963-1965. Robert B. Wassall. March 1966.
ERTM	8	Forest-Fire Pollution Episode in West Virginia, November 8-12, 1964. Robert O. Weedfall. April 1966.
ERTM	9	The Utilization of Radar in Meso-Scale Synoptic Analysis and Forecasting. Jerry D. Hill. March 1966.
ERTM	10	Preliminary Evaluation of Probability of Precipitation Experiment. Carlos R. Dunn. May 1966.
ERTM	11	Final Report. A Comparative Verification of 300 mb. Winds and Temperatures Based on NMC Computer Products Before and After Manual Processing. Silvio G. Simplicio. May 1966.
ERTM	12	Summary of Scientific Services Division Development Work in Sub-Synoptic Scale Analysis and Prediction - Fiscal Year 1966. Fred L. Zuckerberg. May 1966.
ERTM	13	A Survey of the Role of Non-Adiabatic Heating and Cooling in Relation of the Development of Mid-Latitude Synoptic Systems. Constantine Zois. July 1966.
ERTM	14	The Forecasting of Extratropical Onshore Gales at the Virginia Capes. Glen V. Sachse. August 1966.
ERTM	15	Solar Radiation and Clover Temperatures. Alex J. Kish. September 1966.
ERTM	16	The Effects of Dams, Reservoirs and Levees on River Forecasting. Richard M. Greening. September 1966.
ERTM	17	Use of Reflectivity Measurements and Reflectivity Profiles for Determining Severe Storms. Robert E. Hamilton. October 1966.
ERTM	18	Procedure for Developing a Nomograph for Use in Forecasting Phenological Events from Growing Degree Days. John C. Purvis and Milton Brown. December 1966.
ERTM	19	Snowfall Statistics for Williamsport, Pa. Jack Hummel. January 1967
ERTM	20	Forecasting Maturity Date of Snap Beans in South Carolina. Alex J. Kish. March 1967.
ERTM	21	New England Coastal Fog. Richard Fay. April 1967.
ERTM	22	Rainfall Probability at Five Stations Near Pickens, South Carolina, 1957-1963. John C. Purvis. April 1967.
WBTM ER	23	A Study of the Effect of Sea Surface Temperature on the Areal Distribution of Radar Detected Precipitation Over the South Carolina Coastal Waters. Edward Paquet. June 1967. (PB-180-612).
WBTM ER	24	An Example of Radar as a Tool in Forecasting Tidal Flooding. Edward P. Johnson. August 1967 (PB-180-613).
WBTM ER	25	Average Mixing Depths and Transport Wind Speeds over Eastern United States in 1965. Marvin E. Miller. August 1967. (PB-180-614).
WBTM ER	26	The Sleet Bright Band. Donald Marier. October 1967. (PB-180-615).
WBTM ER	27	A Study of Areas of Maximum Echo Tops in the Washington, D.C. Area During the Spring and Fall Months. Marie D. Fellechner. April 1968. (PB-179-339).
WBTM ER	28	Washington Metropolitan Area Precipitation and Temperature Patterns. C.A. Woollum and N.L. Canfield. June 1968. (PB-179-340).
WBTM ER	29	Climatological Regime of Rainfall Associated with Hurricanes after Landfall. Robert W. Schoner. June 1968. (PB-179-341).
WBTM ER	30	Monthly Precipitation - Amount Probabilities for Selected Stations in Virginia. M.H. Bailey. June 1968. (PB-179-342).
WBTM ER	31	A Study of the Areal Distribution of Radar Detected Precipitation at Charleston, S.C. S.K. Parrish and M.A. Lopez. October 1968. (PB-180-480).
WBTM ER	32	The Meteorological and Hydrological Aspects of the May 1968 New Jersey Floods. Albert S. Kachic and William Long. February 1969. (Revised July 1970). (PB-194-222).
WBTM ER	33	A Climatology of Weather that Affects Prescribed Burning Operations at Columbia, South Carolina. S.E. Wasserman and J.D. Kanupp. December 1968. (COM-71-00194).
WBTM ER	34	A Review of Use of Radar in Detection of Tornadoes and Hail. R.E. Hamilton. December 1969. (PB-188-315).
WBTM ER	35	Objective Forecasts of Precipitation Using PE Model Output. Stanley E. Wasserman. July 1970. (PB-193-378).
WBTM ER	36	Summary of Radar Echoes in 1967 Near Buffalo, N.Y. Richard K. Sheffield. September 1970. (COM-71-00310).
WBTM ER	37	Objective Mesoscale Temperature Forecasts. Joseph P. Sobel. September 1970. (COM-71-0074).

**NOAA Technical Memoranda NWS**

NWS	ER 38	Use of Primitive Equation Model Output to Forecast Winter Precipitation in the Northeast Coastal Sections of the United States. Stanley E. Wasserman and Harvey Rosenblum. December 1970. (COM-71-00138).
NWS	ER 39	A Preliminary Climatology of Air Quality in Ohio. Marvin E. Miller. January 1971. (COM-71-00204).
NWS	ER 40	Use of Detailed Radar Intensity Data in Mesoscale Surface Analysis. Robert E. Hamilton. March 1971. (COM-71-00573).
NWS	ER 41	A Relationship Between Snow Accumulation and Snow Intensity as Determined from Visibility. Stanley E. Wasserman and Daniel J. Monte. (COM-71-00763). January 1971.
NWS	ER 42	A Case Study of Radar Determined Rainfall as Compared to Rain Gage Measurements. Martin Ross. July 1971. (COM-71-00897).
NWS	ER 43	Snow Squalls in the Lee of Lake Erie and Lake Ontario. Jerry D. Hill. August 1971. (COM-72-00959).
NWS	ER 44	Forecasting Precipitation Type at Greer, South Carolina. John C. Purvis. December 1971. (COM-72-10332).
NWS	ER 45	Forecasting Type of Precipitation. Stanley E. Wasserman. January 1972. (COM-72-10316).

(CONTINUED ON INSIDE REAR COVER)

**NOAA Technical Memorandum NWS ER-105**

**DENSE FOG CLIMATOLOGY FOR THE BLUE RIDGE FOOTHILLS AND  
PIEDMONT AREAS OF THE BLACKSBURG, VA COUNTY WARNING  
AREA FOR THE PERIOD 1973 -2008**

**JAN JACKSON, KEN KOSTURA AND WILLIAM PERRY**  
National Weather Service Office  
BLACKSBURG, VA

Scientific Services Division  
Eastern Region Headquarters  
Bohemia, New York  
February 2011

United States  
Department of Commerce  
Gary Locke  
Secretary

National Oceanic and  
Atmospheric Administration  
Jane Lubchenco  
Under Secretary and Administrator

National Weather Service  
John L. Hayes  
Assistant Administrator



# Table of Contents

<a href="#">Abstract</a> .....	1
1. <a href="#">Introduction</a> .....	1
2. <a href="#">Data</a> .....	2
3. <a href="#">Methodology</a> .....	2
4. <a href="#">Categories</a> .....	3
4.1 <a href="#">Advection Fog</a> .....	4
4.2 <a href="#">Frontal/Precip. Fog</a> .....	5
4.3 <a href="#">Radiation Fog</a> .....	6
4.4 <a href="#">Radiation Fog/No Front</a> .....	7
4.5 <a href="#">Radiation Fog/Post Cold Front</a> .....	8
4.6 <a href="#">Radiation Fog/Post Warm Front</a> .....	9
5. <a href="#">Discussion</a> .....	10
<a href="#">Acknowledgements</a> .....	11
<a href="#">References</a> .....	11
<a href="#">Figures</a> .....	13
<a href="#">Appendix</a> .....	35

## ABSTRACT

WFO Blacksburg issues Terminal Aerodrome Forecasts (TAFs) for five airports. Dense fog with visibilities of a half mile or less producing VLIFR conditions is a frequent forecast problem at individual airports. In addition, dense fog occasionally becomes widespread enough to necessitate a Dense Fog Advisory. Historical dense fog data is available for each of the five airports. The locations of three of the airports, Roanoke, Lynchburg, and Danville, represent a climatological region with similar geographical characteristics on the east side of the Appalachian Mountains. Thirty six years of fog data (1973 through 2008) from these airports were studied to produce a climatology of dense fog events. Although a quarter of a mile or less visibility is the threshold for a dense fog advisory, we found that we needed to use visibilities of a half mile or less in order to get a large enough sample for the study. A dense fog event was then defined as one in which 2 of the 3 airports recorded dense fog of a half mile or less for at least 2 hours during the same period. A period of two hours or longer would have a major impact on aviation and ground travel, and with the similar topography and relative close proximity of the three airports, the dense fog was then probably extensive enough to warrant a dense fog advisory. 176 separate events were identified in the 36 year period.

---

## 1. INTRODUCTION

The purpose of this study is to develop a climatology of dense fog for the foothills and piedmont region of Blacksburg's County Warning Area (CWA), and use it to help recognize synoptic patterns that are conducive to the formation of dense fog. The foothill and piedmont region was separated from the remainder of the CWA because the topography of the CWA is characterized by two separate regions. There is mainly mountainous terrain over the northwestern half of the CWA along and west of the Blue Ridge mountain range, with some elevated valleys. The mean annual number of days with dense fog in the Blacksburg CWA is highest west of the Blue Ridge Mountains, at greater than 50 days, but an axis of greater than 40 days extends into the foothills and piedmont of southwest Virginia and

northwest North Carolina ([Peace Jr. 1968](#)). The southeastern CWA is comprised of the foothills and piedmont, east of the Blue Ridge Mountains. There is a gradual increase in elevation from southeast to northwest from the piedmont to the Blue Ridge foothills, and then a rapid increase in elevation further northwest into the mountains. Elevations start at around 500 feet in the Piedmont, rise to around 1000 feet in the foothills, and then increase rapidly to mountainous terrain of 2500 to 5000 feet in the Blue Ridge and Appalachian Mountains of southwestern Virginia (VA), southeastern West Virginia (WV), and northwestern North Carolina (NC). In addition, the availability of historical fog data across the CWA was limited to first order airports as defined by the National Climatic Data Center. There are three airports that were judged to be in close enough proximity and elevation

to be representative of the area; Roanoke, Lynchburg and Danville ([Fig. 1](#)).

The total economic losses associated with the impact of the presence of fog on aviation, marine and land transportation can be comparable to that of other severe weather like tornadoes, winter storms and hurricanes. ([NOAA Economics 2008](#)) This local dense fog climatological study will provide forecasters an improved understanding of the frequency and type of dense fog events in the piedmonts of southwest Virginia and northwest North Carolina. ([Croft et al. 1997](#)) This is expected to help forecasters in two areas. The first is for Terminal Aerodrome Forecasts (TAFs) at Roanoke, Lynchburg and Danville. Dense fog would bring either Low Instrument Flight Rules (LIFR) of visibilities less than a mile, or Very Low Instrument Flight Rules (VLIFR) of visibilities less than a half mile ([NWSI 10-813](#) November 19, 2007). The second area this study is expected to help forecasters with is the issuance of dense fog advisories for widespread fog with visibilities of a quarter of a mile or less ([NWSI 10-515](#) NOVEMBER 1, 2005).

## 2. DATA

Data were gathered for all dense fog events for the airports of Roanoke, Lynchburg and Danville during the period of January 1973 through December 2008. The climate database in the TAF monitor program, Aviation Forecast Preparation System (AvnFPS), was used to find the periods of dense fog at each site. For the purposes of this study, one half mile visibility or less was used to define dense fog. There were two reasons for this. First, a half mile or

less corresponds closely to VLIFR conditions. Second, although a quarter of a mile or less visibility is the threshold for a dense fog advisory, we found that we needed to use visibilities of a half mile or less in order to get a large enough sample for the study. A dense fog event was then defined as one in which 2 of the 3 sites recorded dense fog of a 1/2 mile or less for at least 2 hours during the same period. A period of two hours or longer would have a major impact on aviation and ground travel, and with the similar topography and relative close proximity of the three airports, the dense fog was then probably extensive enough to warrant a dense fog advisory. 176 separate events were identified in the 36 year period.

There was a wide range in the annual distribution of dense fog events. In four of the 36 years of the study there were no dense fog events at all, and nearly half (17) of the years had three events or less. In comparison, eight of the years had 10 or more dense fog events, with a maximum of 18 events identified in 1982. Dense fog events were found in every month. The distribution by month is characterized by a minimum in the spring and summer months (just 25 events in the April through August period), then a sharp increase in the number of events to a peak in November, with high numbers through the winter months (December, January, February) when there is a minimum of solar radiation. ([Slemmer 2004](#); [Fig. 2a](#)).

## 3. METHODOLOGY

There are three types of fog that were identified from the dense fog events. The fog types were determined

using the characteristics comparison table from the COMET<sup>®</sup> Distance Learning Aviation Course (DLAC) on fog, ([Bol 2001](#)), and reviewing the surface and upper air data, including soundings at Greensboro and Blacksburg (available at Blacksburg only after 1995).

Synoptic and upper air patterns, along with the presence and timing of any precipitation, were utilized to help classify the categories of dense fog events. Details of the start and end times of the fog events and whether there was precipitation with the fog can be found in the [Appendix](#).

The types of fog identified in this study are radiation (or ground) fog, advection fog, and frontal/precipitation fog. Further, for radiation fog, three distinct synoptic patterns were determined based on review of surface data and upper air patterns, and the timing of synoptic frontal features with any associated precipitation. Dense radiation fog events that developed within 24 hours of a warm frontal passage were categorized as Post Warm Front, and events that developed within 12 hours of a cold frontal passage were categorized as Post Cold Front. Dense radiation fog events under synoptic high pressure (and no frontal passage within 24 hours) were labeled as No Front events.

Therefore, for the purpose of this study, we have five categories of dense fog events. The categories are: Advection Fog (23 events), Frontal/Precipitation Fog (41 events), Radiation Fog/No Front (39 events), Radiation Fog/Post Cold Front (34 events), and Radiation Fog/Post Warm

Front (39 events) ([Fig. 2b](#)). The distribution of fog type by month shows that Advection, Frontal/Precipitation, and Radiation/Post Warm Front fog all occurred only in the cooler months, (September through April), while Radiation/Post Cold Front and Radiation/No Front occurred in the warm season, (May through August), as well as the cooler months ([Fig. 2c](#)).

In an effort to show the synoptic pattern of each category, composite maps of various meteorological fields were generated using the Earth System Research Laboratory (ESRL) 6-Hourly Composites web page which uses the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) reanalysis data. This was accomplished by assigning an actual start, middle and ending time of the dense fog in each of the cases in a category to the nearest 6-hour synoptic time, (00, 06, 12, 18 UTC). The maps generated then represented a composite for a particular field around the start, middle and end times of the dense fog for each category.

## 4. CATEGORIES

### 4.1. ADVECTION FOG (23 events)

Advection fog develops when warm, moist air moves over a colder underlying surface. The surface could be cold ground, snow cover, water or ice. The fog forms primarily by boundary layer dynamic and adiabatic processes including advection of moisture and temperature. It often occurs with light or moderate low-level winds (up through 850 mb) that are less than  $10 \text{ ms}^{-1}$ , but can also occur with winds stronger than  $10 \text{ ms}^{-1}$ . Rapid development can occur

as the preconditioned air mass moves over a cooler surface.

Low-level factors to look for with advection fog include: air parcel trajectories originating over a moisture source sufficient to establish a moist boundary layer condition, the depth of the surface-based moist layer increases as a result of mechanical turbulence and convective mixing of buoyant moist air, and cold air damming with large-scale anticyclonic winds and subsidence providing a capping inversion so the boundary layer can saturate.

In identifying the 23 advection fog events, Greensboro NC (GSO) soundings were utilized to determine the low level temperature and dewpoint profiles, and the presence of cold air damming. The low-level winds in the soundings helped identify warm advection. Historical daily weather maps provided by the NOAA Central Library Data Imaging Project were utilized to determine the presence and location of any high or low pressure centers, and synoptic scale fronts. The surface observations at all three of the sites were used to determine if there was precipitation during the event. Many of the advection fog cases had drizzle reported during the event. Five of the advection fog events had light rain reported at one or more of the sites, but the daily weather maps indicated there was no warm front near the region.

Advection fog can last several days and may be advected over large areas and across great distances. The average length of the 23 advection fog events was 13.6 hours, the longest of all the fog categories. It tends to be deeper than radiation fog since it is often driven by

synoptic scale factors, and it can form and advect into a region almost any time of day, although it has some tendency to develop in the evening hours as the boundary layer cools.

#### a) Synoptic Overview

Composite maps of mean sea level pressure were generated for the start, middle and end of the advection fog events ([Fig. 3a](#)). A start time was determined for each event by assigning the nearest 6 hour synoptic time (00, 06, 12, 18 UTC), in which one or more of the airports first reported dense fog. The ending time for each event was determined by assigning the nearest 6 hour synoptic time to the last observation of dense fog at any of the three airports. The middle point of each event was then assigned to the nearest 6 hour synoptic time to the average between the start and end times. The surface map at the start shows a high pressure centered off the mid-Atlantic coast, with low pressure over the western Great Lakes, a warm front across the Ohio Valley and into the southern Appalachians, and a cold front in the Mississippi Valley. By the middle of the advection fog events, the center of the high pressure was further east into the Atlantic, while the low pressure center and trailing cold front were moving slowly eastward. The warm front was north of the study area. By the end of the advection fog events, the high pressure was well east of the region and the cold front had progressed into the Tennessee Valley.

Composite maps of 925 mb relative humidity, 850 mb relative humidity, and surface winds for the day of the event ([Figs. 3b](#), [3c](#) and [3d](#)) indicate shallow



low-level moisture over the study area, with near saturation to 850 mb, and very little change in the low-level moisture profile through the event. There were east to southeast surface winds of less than  $4 \text{ ms}^{-1}$  into the study area, with trajectories off the Atlantic. Winds above the boundary layer at 850 mb (not shown) were 8 to  $12 \text{ ms}^{-1}$  from the southwest, indicating warm advection. The upper pattern in the advection fog cases (not shown) had a progressive, full latitude Midwest trough, and a southeast U.S. upper ridge.

Representative GSO soundings for advection fog from November 30, 1977 (Fig. 4) show uniform southwest flow (warm advection) above a shallow saturated layer through the event. In this case, dense fog developed at Roanoke and Lynchburg around 30/00 UTC, developed at Danville at 29/14 UTC, and lasted until 29/18 UTC for all sites. The only precipitation reported was light drizzle.

There were 23 events identified as advection fog. This represents just 13% of the total, making advection fog the rarest form of widespread dense fog. Indeed, over half (19) of the years in the study period had no cases of dense advection fog. Advection fog occurred only in the cooler months (September through March), with a peak from November through January.

#### *4.2. FRONTAL/PRECIPITATION FOG* (41 events)

Frontal/precipitation fog is formed by precipitation falling into a cold stable layer and raising the dewpoint. The fog is formed primarily by diabatic processes - evaporational

cooling/condensation. The evaporation/condensation process extends the cloud to the surface. The fog may develop during the precipitation, or after the precipitation ends. It is most common with cold air damming in the vicinity of warm or stationary fronts.

In identifying the 41 frontal/precipitation fog events, the surface observations at all three of the sites were used to identify the events with prolonged rainfall. All of the events identified as frontal/precipitation had rain or drizzle reported at each site that had dense fog. Historical daily weather maps were utilized to determine the presence and location of any high or low pressure centers, and synoptic scale fronts. In all of the frontal/precipitation events, a warm front was in the vicinity of the study area during the event. Greensboro NC (GSO) soundings were utilized to look for a deep saturated layer with a veering wind profile indicating warm advection and overrunning. The main factors in differentiating between frontal/precipitation fog and advection fog were whether or not there was precipitation, and how long it lasted, along with the presence of a warm front.

Frontal/precipitation fog may develop over large areas and across great distances, and can last a day or two. The average length of the 41 Frontal/Precipitation Fog events was 10.1 hours, the second longest of all the fog categories. The depth of the fog is normally greater than any other type. It can form any time of the day or night.

### a) Synoptic Overview

Composite maps of mean sea level pressure were generated for the start, middle and end of the frontal/precipitation fog events (Fig. 5a). The surface map at the start shows a warm front extending eastward from a low pressure in the Ohio Valley, and across the area of study. A trailing cold front extends southward through the Tennessee Valley. By the middle of the frontal/precipitation fog events, the warm front had moved north of the area of study into northern Virginia, while the cold front had progressed eastward but remained west of the Appalachians. All of the area of study is in the warm sector. At the end of the frontal/precipitation fog events, the cold front had moved into the mountains just west of the study area.

Composite maps of 925 mb relative humidity, 850 mb relative humidity, and surface winds for the day of the event (Figs. 5b, 5c and 5d) indicate near saturation to 850 mb over the study area, with very little change through the event. Surface winds are light, generally less than  $3 \text{ ms}^{-1}$ , while 850 mb winds (not shown) are from the west-southwest at 11 to  $14 \text{ ms}^{-1}$ . The upper level pattern associated with frontal/precipitation fog (not shown) features a progressive, full latitude Midwest trough, with a strong Great Lakes short wave, and a weak southeast U.S. upper ridge.

Representative GSO soundings for frontal/precipitation fog from December 2, 1991 (Fig. 6) show a strong, slightly veering, south to southwest flow (warm advection) and a deep nearly saturated layer from the surface to 750 mb at the beginning of the event (02/00 UTC). By

the middle of the event (02/12 UTC), the saturated layer extends from the surface to 600 mb. In this case, rain began before 01/12 UTC at Roanoke and Lynchburg, and the ceilings and visibility lowered to 0.5 mile or less and 100 feet after 02/00 UTC, then rose some during the day (after 02/12 UTC), before dropping back again after 03/00 UTC with more rain.

Frontal/precipitation events with widespread dense fog were more common than advection fog in the area of study. Forty one of the 176 events (22%) were from frontal/precipitation fog. Even though it was more common than advection fog, there were still 13 years in the study period with no frontal/precipitation fog events identified. The monthly distribution of frontal/precipitation fog was similar to advection fog, only occurring in the cooler months, (September through April), with a peak from November through February.

### 4.3. RADIATION FOG (112 events)

Radiative cooling is the most common process by which dense fog is formed over the study area. Radiation fog is produced over a land area when radiational cooling reduces the air temperature to its dewpoint. There were three types of radiation fog that affected the study area, based upon their distinct synoptic pattern.

One of the types, Radiation Fog/No Front, (RF/No), occurred under the influence of high pressure, with no frontal passage and no rainfall within 24 hours of the event. A ridge of high pressure was located over the southeastern U.S., and low level winds

were light and from the southeast to south.

A second type, Radiation Fog/Post Cold Front, (RF/Cold), occurred within 12 hours after a weak cold front or surface trough passage. In these events, dense fog occurred with surface high pressure building into the region from the west with low level winds veering from the south before the event, to the southwest or west near the beginning of the event. GSO soundings indicated a moist layer below 700 mb rapidly dried out and radiative cooling left a very shallow saturated layer at the surface.

A third type, Radiation Fog/Post Warm Front, (RF/Warm), occurred within 24 hours after a warm frontal passage. Most of the RF/Warm events had rainfall reported *before* the dense fog developed, but no rainfall reported *during* the dense fog. GSO soundings were utilized to determine the progression and depth of the saturated layer, and all of the events identified as RF/Warm indicated a nearly saturated layer up to 700 – 500 mb, drying in the mid-levels before dense fog developed. Although the rainfall moistened and pre-conditioned the lower levels of the atmosphere, radiative cooling was determined to be the process by which dense fog developed.

All three types of radiation fog occur in the presence of a weak low-level anti cyclone with moist surface conditions under a capping inversion, and moderate vertical shear near the capping inversion. There is rapid nocturnal radiational cooling of the lower boundary layer.

Radiation fog tends to be short duration, less than 24 hours. The

average length of the 112 radiation fog events was 8 hours. RF/No and RF/Warm both averaged around 8.5 hours, while RF/Cold had the shortest average length of all dense fog categories at 7.2 hours. Radiation fog tends to be confined to a shallow layer that mainly forms late at night or early in the morning. Surface winds are generally  $5 \text{ ms}^{-1}$  or less.

When considering all three types, this general category is by far the most common producer of widespread dense fog in the study area. There were 112 radiation fog events, representing 64% of the total. The three types of radiation fog and a description of their synoptic patterns follows.

#### 4.4 RADIATION FOG/NO FRONT (39 events)

##### a) Synoptic Overview

Composite maps of mean sea level pressure were generated for the start, middle, and end of the RF/No events ([Fig. 7a](#)). The surface map at the start the RF/No events shows a large high pressure centered off the New England coast, and strong ridging into the southeastern U.S. A cold front extends from the western Great Lakes, southwestward into Texas. A warm front extends eastward into Pennsylvania from the low pressure. By the middle of the event, surface high pressure remains ridged strongly into the southeast U.S., while the surface low pressure and associated cold front has progressed slowly eastward. The RF/No events end as the low pressure and cold front move into the Ohio and Tennessee Valleys, and the surface high pressure over the southeast U.S. weakens.

Composite maps of 925 mb relative humidity, 850 mb relative humidity, and surface winds for the day of the event (Figs 7b, 7c, and 7d) indicate very shallow low level moisture, with greater than 80 percent saturation at 925 mb and greater than 70 percent saturation at 850 mb through the event. The 850 and 925 mb levels became increasingly saturated west of the Appalachians during the event. Surface winds are less than  $4 \text{ ms}^{-1}$  from the southwest, and 850 mb winds (not shown) are 8 to  $12 \text{ ms}^{-1}$  from the west-southwest. The upper pattern in RF/No events features a weakening Midwest trough, with a short wave lifting through the Great Lakes region.

Representative soundings from December 29, 1974 for RF/No (Fig. 8) show a shallow saturated layer near the beginning of the event at 29/00UTC, that transitions to a saturated layer from the surface to 850 mb by 29/12 UTC. There were light northeast surface winds near the beginning of the event at 29/00 UTC. Surface winds by 29/12 UTC were light despite a strengthening west flow above the boundary layer. In this case, dense fog began around 29/02 UTC, ended at 29/1930 UTC. Dense fog was reported at all three sites. There was no precipitation during the period.

Thirty nine of the 176 dense fog events were identified as RF/No. RF/No occurred in every month, except for February, with the peak months (27 events) extending from August through December. Similar to all radiation fog types, nearly half (17) of the years in the study had no widespread dense fog events identified as RF/No.

#### 4.5 RADIATION FOG/POST COLD FRONT (34 events)

##### a) Synoptic Overview

Composite maps of mean sea level pressure were generated for the start, middle and end of the radiation fog/post cold front events (Fig. 9a). The surface map at the start of the RF/Cold events shows a weak low pressure over the Maryland/Delaware region, with a trailing cold front off of the mid-Atlantic coast. Weak high pressure is over the Gulf Coast region. In the middle of the event, the low pressure and cold front move further off shore, while the high pressure builds northward into the Tennessee valley. The RF/Cold events end when the high pressure is centered over the southern Appalachians.

Composite maps of 925 mb relative humidity, 850 mb relative humidity, and surface winds for the day of the event (Figs 9b, 9c, and 9d) indicate near saturated low levels at the start of the event, (greater than 90 percent at 925 mb and 80 percent at 850 mb), gradually drying out from the top down through the middle and into the end of the event with the relative humidity dropping to 60 percent at 850 mb near the end of the event. There is a light southwest surface wind of less than  $3 \text{ ms}^{-1}$ , and west-southwest 850 mb winds (not shown) of 7 to  $8 \text{ ms}^{-1}$ . The upper level pattern associated with RF/Cold events (not shown) features a weak short wave passing across the region in southwest flow aloft.

Representative soundings for RF/Cold from November 7, 1989 (Fig. 10) show a moist layer below 850 mb near the beginning of the event (07/00

UTC). There was drying above 950 mb, a shallow saturated layer under a steep inversion near the surface, and lighter surface winds near the peak of the event (07/12 UTC). The main change in airmass behind the front was drier air. In this case, dense fog began shortly after 07/00 UTC at Roanoke and Lynchburg, and after 07/03 UTC at Danville. The dense fog ended around 07/14 UTC at Roanoke, and 07/16 UTC at Lynchburg and Danville. Precipitation preceded the frontal passage at Lynchburg and Danville, but not Roanoke.

Thirty four of the 176 dense fog events were identified as radiation/post cold frontal passage. Radiation fog/post cold frontal passage fog occurred in every month, with minimums in the spring and summer months, (March through August), and a peak in the winter months (December through February). Similar to all radiation fog types, nearly half (16) of the years in the study had no widespread dense fog events identified as RF/Cold type fog.

#### *4.6 RADIATION FOG/POST WARM FRONT (39 events)*

##### *a) Synoptic Overview*

Composite maps of mean sea level pressure were generated for the start, middle and end of the RF/Warm events ([Fig. 11a](#)). At the start of the RF/Warm events, a warm front extends across northern Virginia just north of the area of study, and extends westward to a low pressure over Wisconsin. A cold front extends southward through the Mississippi valley. By the middle of the event, the low pressure has progressed eastward into northwest Michigan, with the trailing cold front into western

sections of the Ohio and Tennessee Valleys. The warm front moved further north of the area of study. The RF/Warm events end when the cold front moves into the mountains of southwest Virginia.

Composite maps of 925 mb relative humidity, 850 mb relative humidity, and surface winds for the day of the event ([Figs 11b](#), [11c](#) and [11d](#)) indicate near-saturated low levels at the start of the event, (greater than 80 percent at 925 and 850 mb, gradually drying out from the top down through the middle and into the end of the event, with the relative humidity dropping to 65 percent at 850 mb near the end of the event. There are light south surface winds less than  $4 \text{ ms}^{-1}$ , and west-southwest 850 mb winds (not shown) of 7 to  $8 \text{ ms}^{-1}$ . The upper level pattern associated with RF/Warm events (not shown) features a full latitude Midwest trough, with southwest flow aloft over the mid-Atlantic region.

Representative soundings for RF/Warm from December 2, 1982 ([Fig. 12](#)) show drying levels above 900 mb near the beginning of the event, (02/00 UTC), and light winds with a steep saturated surface inversion at the peak of the event (02/12 UTC). There was rain at Roanoke and Lynchburg before the dense fog developed at all three sites in this event. Dense fog began as early as 02/00 UTC at Roanoke, and ended as late as 02/1730 UTC at Lynchburg.

Thirty nine of the 176 dense fog events were identified as the RF/Warm fog type. The monthly distribution of RF/Warm was similar to advection fog and frontal/precipitation fog: only occurring in the cooler months, with a peak from October through December.



Similar to all radiation fog types, nearly half (16) of the years in the study had no widespread dense fog events identified as RF/Warm type fog.

## 5. DISCUSSION

Although there were 176 dense fog events identified in a 36 year period, there was a wide variability in the annual occurrence, with no widespread dense fog events at all in four of the 36 years, (1985, 1987, 1988, 1992). In the four null years, three of them had moderate drought conditions on the annual Palmer Drought Severity Index, (PDSI) and one was neutral (NCDC 2008). Since soil moisture can change rapidly from month to month, a look at the monthly historical PDSI showed a range from one to three months in each of the null years with moderate to severe drought (NCDC 2010). More importantly, there were no months during the null years with moist conditions. In comparison, of the eight years with ten or more events, six (1973, 1974, 1975, 1978 and 1979) had several months with moist conditions and no drought. One, (1982 with 18 events), had no drought but only one moist month, and one (2006) had 4 months with moderate drought, but also 4 moist months (NCDC 2010). Further study of soil moisture and other environmental variables, such as percent of normal precipitation, may provide more indicators for widespread dense fog development. There were an average of about five events per year, and a median of four events, indicating that widespread dense fog events in the foothills and piedmont of southwest Virginia and northwest North Carolina are relatively rare. Widespread dense fog is also mostly a cool season phenomena. There were only twenty

three (13%) widespread dense fog events identified in the spring and summer, (April through August time frame). All but one of the dense fog events from April through August were radiation fog. Nearly two thirds of all dense fog events, (115 of 176 or 65%), occurred during November through February, and over half of the events, (92 of 176 or 52%), occurred in the three month November through January period.

All-category composite maps (Figs 13, 14, 15, 16, 17, 18, 19) combined all categories at the middle of the events. These all-category composites showed that a key feature to look for in the surface synoptic pattern is a surface high pressure centered along or off the east coast (Fig. 13). This is the case in all but the RF/Cold type. However, the location of any warm frontal boundaries becomes less distinct in the all category composite. In a majority of the dense fog cases, (frontal/precipitation and RF/Warm combined), a surface warm front will be in close proximity to the study area. The composite maps from all categories also show a warm advection pattern, with veering winds from the surface to 850 mb (Figs. 14 and 15). This veering wind pattern, from southeast to south at the surface, to southwest at 850 mb, can be found in all of the individual category composites except for RF/Cold. Even though with the RF/Cold type there is weak cold advection, the relatively small number of this type means that signal is averaged out in the composites. Other key features to note at the surface from the all-category composites are that surface winds are light,  $4 \text{ ms}^{-1}$  or less (Fig. 14), but not necessarily calm. All-category composite surface winds are from a southerly direction. 850 mb winds are

from the southwest, with the core of the strongest winds in either the Tennessee or Ohio Valleys, extending across the central Appalachians (Fig. 15). Wind speeds at 850 mb over the study area are around  $10 \text{ ms}^{-1}$ .

The synoptic pattern in the upper levels is similar in all of the dense fog categories, with a central U.S. trough and weak southeast U.S. ridge at 500 mb (Fig. 16). Otherwise, there are only subtle differences between the categories.

As for low level moisture, the composite of all categories at the middle of the event shows relative humidity of 90 percent at 925 mb across the study area (Fig. 17). The 850 mb relative humidity is greater than 70 percent (Fig. 18). The frontal/precipitation and advection fog categories both contribute greater than 80 percent relative humidity at 850 mb, while the radiation fog types are less than 70 percent. The all-category composite 700 mb relative humidity indicates drier air aloft, with only 40 to 45 percent relative humidity

over the study area (Fig. 19). This drier air applies to all categories except for the frontal/precipitation fog (not shown).

In conclusion, sensitivity to the time of year and composites of the patterns that produce the different categories of dense fog events in the Blue Ridge foothills and piedmont areas of Blacksburg, Virginia CWA will result in a better anticipation of dense fog development.

## ACKNOWLEDGEMENTS

The authors would like to thank Steve Keighton (Science and Operations Officer at WFO Blacksburg), Dave Radell (Eastern Region Scientific Services Division), and two anonymous reviewers in the Eastern Region, for their insight and review of the document. Also special thanks to Paul Jendrowski (Information Technology Officer at WFO Blacksburg) for help in developing the topographical map.

## REFERENCES

- Bol, Alan, 2001: COMET: Anticipating Hazardous Weather and Community Risk. [Available online from: <http://www.meted.ucar.edu/hazwx/topic3/fact12.htm>]
- Croft, P.J., R.L. Pfost, J.M. Medlin, and G.A. Johnson, 1997: Fog forecasting for the Southern Region: A conceptual model approach. *Wea. Forecasting*, **12**, 545-556.
- National Weather Service Instruction 10-813, 2007: Terminal Aerodrome Forecasts. [Available online from <http://www.nws.noaa.gov/directives/sym/pd01008013curr.pdf>]
- National Weather Service Instruction 10-515, 2005: WFO Non-Precipitation Weather Products Specification. [Available online from <http://www.nws.noaa.gov/directives/sym/pd01005015curr.pdf>]

NCDC, 2010: Historical Palmer Drought Indices. [Available online at: <http://www.ncdc.noaa.gov/temp-and-precip/drought/historical-palmers.php>

NCDC, 2008: Tree-ring reconstructions of palmer drought severity index across North America over the last 2000 years. [Available online at: <http://www.ncdc.noaa.gov/cgi-bin/paleo/pd04plot.pl>]

NOAA Economics 2008: The Economics and Social Benefits of NOAA Data & Products. [Available Online at: <http://www.economics.noaa.gov/?goal=commerce&file=events/fog>]

Peace, Jr., 1968: Heavy fog regions in the Conterminous United States. *Mon. Wea. Rev.*, **97**, 116-123.

Slemmer, J., 2004: Study of dense fog at the Salt Lake City International Airport and its impacts to aviation. *National Weather Service Technical Attachment*, 2004-01. [Available online from <http://www.wrh.noaa.gov/wrh/04TAs/0401/SLCFog.htm>]

## **DISCLAIMER**

Reference to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not constitute or imply its recommendation, or favoring by the United States Government or NOAA/National Weather Service. Use of information from this publication shall not be used for advertising or product endorsement purposes.



## FIGURES




Figure 1. TAF sites KROA, KLYH and KDAN. White lines indicate the Blue Ridge Mountains.

### Number of Events of Dense Fog Per Month




Figure 2a. Number of dense fog events per month over the study period.




Figure 2b. Number of dense fog events by category for the period of study.

### Monthly Distribution of Fog Types




Figure 2c. Monthly distribution of fog type for the period of study.




Figure 3a. Composite surface map of mean sea level pressure (mb) and overview of advection fog cases. From left to right: Start, middle, and end of event.






Figure 3b. Composite 925 mb relative humidity (%) overview of advection fog. From left to right: Start, middle and end of the event.




Figure 3c. Composite 850 mb relative humidity (%) of advection fog. From left to right: Start, middle and end of the event.




Figure 3d. Composite surface winds (ms<sup>-1</sup>) of advection fog. From left to right: Start, middle and end of the event.




Figure 4. Advection Fog representative soundings at the (a) beginning and (b) during the peak, of the event. Images courtesy of Plymouth State Weather: (<http://vortex.plymouth.edu/>)




Figure 5a. Composite surface map of mean sea level pressure (mb) and overview of frontal/precipitation fog cases. From left to right: Start, middle, and end of event.




Figure 5b. Composite 925 mb relative humidity (%) overview of frontal/precipitation fog. From left to right: Start, middle and end of the event.




Figure 5c. Composite 850 mb relative humidity (%) overview of frontal/precipitation fog. From left to right: Start, middle and end of the event.




Figure 5d. Composite surface winds ( $\text{ms}^{-1}$ ) overview of frontal/precipitation fog. From left to right: Start, middle and end of the event.



(a) GSO Sounding 00UTC 2 Dec 91

(b) GSO Sounding 12UTC 2 Dec 91




Figure 6. Frontal/precipitation Fog representative soundings at the (a) beginning, and (b) during the peak, of the event. Images courtesy of Plymouth State Weather: (<http://vortex.plymouth.edu/>)




Figure 7a. Composite surface map of mean sea level pressure (mb) and overview of fog/no front cases. From left to right: Start, middle, and end of event.






Figure 7b. Composite 925 mb relative humidity (%) overview of radiation fog/no front. From left to right: Start, middle and end of the event.




Figure 7c. Composite 850 mb relative humidity (%) overview of radiation fog/no front. From left to right: Start, middle and end of the event.




Figure 7d. Composite surface winds ( $\text{ms}^{-1}$ ) overview of radiation fog/no front. From left to right: Start, middle and end of the event.

(a) GSO Sounding 00UTC 29 Dec 74

(b) GSO Sounding 12UTC 29 Dec 74




Figure 8. Radiation fog/no front representative soundings at the (a) beginning, and (b) during the peak, of the event. Images courtesy of Plymouth State Weather: (<http://vortex.plymouth.edu/>)




Figure 9a. Composite surface map of mean sea level pressure (mb) and overview of radiation fog/post cold front cases. From left to right: Start, middle, and end of event.




Figure 9b. Composite 925 mb relative humidity (%) overview of radiation fog/post cold front. From left to right: Start, middle and end of the event.




Figure 9c. Composite 850 mb relative humidity (%) overview of Radiation fog/post cold front. From left to right: Start, middle and end of the event.





Figure 9d. Composite surface winds ( $\text{ms}^{-1}$ ) overview of Radiation fog/post cold front. From left to right: Start, middle and end of the event.



(a) GSO Sounding 00UTC 7 Nov 89



(b) GSO Sounding 12UTC 7 Nov 89




Figure 10. Radiation fog/post cold front representative soundings at the (a) beginning, and (b) during the peak of the event. Images courtesy of Plymouth State Weather: (<http://vortex.plymouth.edu/>)




Figure 11a. Composite surface map of mean sea level pressure (mb) and overview of radiation fog/post warm front cases. From left to right: Start, middle, and end of event.




Figure 11b. Composite 925 mb relative humidity (%) overview of radiation fog/post warm front. From left to right: Start, middle and end of the event.




Figure 11c. Composite 850 mb relative humidity (%) overview of radiation fog/post warm front. From left to right: Start, middle and end of the event.




Figure 11d. Composite surface wind ( $\text{ms}^{-1}$ ) map overview of radiation fog/post warm front. From left to right: Start, middle and end of the event.




Figure 12. Radiation fog/post warm front representative soundings at the (a) beginning, and (b) during the peak, of the event. Images courtesy of Plymouth State Weather: (<http://vortex.plymouth.edu/>)




Figure 13. Mean sea level pressure (mb) composite for all categories.







Figure 14. Surface wind speed ( $\text{ms}^{-1}$ ) composite for all categories.



850mb Vector Wind (m/s) Composite Mean  
 9/9/74 12/29/74 11/7/96 11/27/01 2/17/75 10/26/64 11/7/69 2/15/05 2/18/75 1/20/62 2/8/86 2/  
 NCEP/NCAR Reanalysis

Figure 15. 850mb wind speed ( $\text{ms}^{-1}$ ) composite for all categories.




Figure 16. 500 mb height (m) composite for all categories.






Figure 17. 925 mb relative humidity (%) composite for all categories.



850mb Relative Humidity (%) Composite Mean  
 9/9/74 12/29/74 11/7/96 11/27/01 2/17/75 10/26/64 11/7/69 2/15/05 2/18/75 1/20/62 2/8/86 2/  
 NCEP/NCAR Reanalysis

Figure 18. 850 mb relative humidity (%) composite for all categories.



700mb Rel. Humidity (%) Composite Mean  
 74 12z 11/7/98 0z 11/28/01 0z 2/17/75 6z 10/27/84 0z 11/8/89 0z 2/15/05 6z 2/18/75 12z 1/20/8  
 NCEP/NCAR Reanalysis

Figure 19. 700 mb relative humidity (%) composite.

## APPENDIX

### Dense Fog Events by Category With Beginning and Ending Times UTC And Precipitation Other Than Drizzle

#### I. Advection Fog (23 events)

Beginning	Ending	Locations with Fog	Precipitation
03/06/73 0700	03/06/73 1400	ROA and DAN	No
10/13/73 0900	10/13/73 1400	ROA and LYH	No
12/25/73 1100	12/25/73 1700	ROA and LYH	No
01/03/74 1700	01/04/74 0300	ROA and LYH	No
01/26/74 1800	01/27/74 0600	ROA and LYH	No
11/19/74 0100	11/19/74 1700	ROA, LYH and DAN	No
11/09/75 0900	11/09/75 1500	LYH and DAN	No
01/25/76 1800	01/26/76 2000	ROA, LYH and DAN	Yes
11/30/77 0000	11/30/77 1900	ROA, LYH and DAN	No
02/17/78 0400	02/17/78 1800	LYH and DAN	No
09/28/79 1200	09/29/79 1300	ROA, LYH and DAN	Yes
02/04/82 0200	02/04/82 1500	ROA, LYH and DAN	No
12/03/82 0400	12/03/82 1800	ROA, LYH and DAN	No
10/12/83 1000	10/12/83 1500	LYH and DAN	No
12/16/84 0900	12/16/84 1200	ROA, LYH and DAN	No
12/29/90 0600	12/30/90 1300	ROA, LYH and DAN	No
01/17/96 0700	01/18/96 1100	ROA, LYH and DAN	No
01/06/98 1200	01/07/98 0600	ROA and LYH	Yes
12/19/02 2300	12/20/02 08	LYH and DAN	Yes
03/01/03 0100	03/01/03 1600	ROA and LYH	No
11/18/03 1800	11/18/03 2300	ROA and LYH	Yes
11/28/05 1600	11/28/05 2300	ROA and DAN	No
12/18/08 0500	12/18/08 2300	ROA, LYH and DAN	No

## II. Frontal/Precipitation Fog (41 events)

Beginning	Ending	Locations with Fog	Precipitation
12/14/03 0000	12/14/03 1000	LYH and DAN	Yes
12/20/73 2100	12/21/73 1000	LYH and DAN	Yes
01/01/74 0400	01/01/74 1100	ROA and LYH	Yes
01/07/76 2100	01/08/76 1300	ROA and LYH	Yes
12/12/76 0000	12/12/76 1500	ROA, LYH and DAN	Yes
09/17/77 0300	09/17/77 1300	ROA and LYH	Yes
01/08/78 1700	01/09/78 0000	LYH and DAN	Yes
01/25/78 2000	01/26/78 0300	LYH and DAN	Yes
04/19/78 1400	04/20/78 0200	LYH and DAN	Yes
11/15/78 2300	11/16/78 2000	ROA, LYH and DAN	Yes
11/17/78 1800	11/18/78 0200	LYH and DAN	Yes
01/07/79 0300	01/07/79 1900	ROA, LYH and DAN	Yes
01/13/79 1500	01/14/79 1400	ROA, LYH and DAN	Yes
01/21/79 0400	01/21/79 1500	LYH and DAN	Yes
02/23/79 2200	02/24/79 1800	ROA and LYH	Yes
01/11/80 0900	01/12/80 0300	ROA, LYH and DAN	Yes
02/20/81 0200	02/20/81 0800	ROA and LYH	Yes
12/27/81 0800	12/27/81 1500	LYH and DAN	Yes
01/03/82 1900	01/04/82 0300	ROA, LYH and DAN	Yes
01/23/82 2100	01/24/82 0100	ROA and LYH	Yes
03/17/82 0000	03/17/82 1100	ROA and LYH	Yes
03/20/82 2330	03/21/82 1600	ROA, LYH and DAN	Yes
12/15/82 2200	12/16/82 0500	ROA, LYH and DAN	Yes
01/23/83 1200	01/23/83 1900	LYH and DAN	Yes
02/06/83 0800	02/06/83 1500	ROA and DAN	Yes
02/04/84 0300	02/04/84 1200	ROA, LYH and DAN	Yes
02/04/86 0600	02/05/86 1100	ROA, LYH and DAN	Yes
12/02/91 0400	12/02/91 1100	ROA, LYH and DAN	Yes
01/13/93 1000	01/13/93 1500	ROA, LYH and DAN	Yes
02/22/93 0500	02/22/93 1300	ROA, LYH and DAN	Yes
02/12/94 2100	02/13/94 1300	ROA, LYH and DAN	Yes
11/28/94 0400	11/28/94 1600	LYH and DAN	Yes
11/02/95 0200	11/02/95 1300	ROA, LYH and DAN	Yes
09/17/96 0200	09/17/96 1000	ROA, LYH and DAN	Yes
12/17/96 0400	12/17/96 1200	ROA, LYH and DAN	Yes
12/04/97 0900	12/04/97 1400	ROA, LYH and DAN	Yes
12/13/99 2300	12/14/99 0600	ROA, LYH and DAN	Yes
01/10/00 0200	01/10/00 1100	ROA, LYH and DAN	Yes
02/04/06 1300	02/04/06 2100	ROA, LYH and DAN	Yes
02/11/06 1800	02/11/06 2200	ROA and LYH	Yes
01/05/07 1300	01/05/07 1800	ROA and LYH	Yes



### III. Radiation No Front Fog (39 cases)

Beginning	Ending	Locations with Fog	Precipitation
03/04/73 0100	03/04/73 1430	ROA and DAN	No
09/11/73 0900	09/11/73 1330	ROA and LYH	No
01/10/74 0800	01/10/74 1500	ROA and DAN	No
08/16/74 0800	08/16/74 1330	LYH and DAN	No
08/27/74 0900	08/27/74 1300	ROA and LYH	No
09/09/74 0500	09/09/74 1400	ROA, LYH and DAN	No
12/29/74 0200	12/29/74 1930	ROA, LYH and DAN	No
10/01/75 1000	10/01/75 1500	ROA and DAN	No
07/29/76 0700	07/29/76 1200	ROA and LYH	No
11/09/77 1000	11/09/77 1600	LYH and DAN	No
11/14/78 1000	11/14/78 1500	LYH and DAN	No
08/26/79 0900	08/26/79 1300	ROA and LYH	No
08/28/79 0500	08/28/79 1300	ROA and LYH	No
09/04/79 0900	09/04/79 1400	LYH and DAN	No
10/02/79 0900	10/02/79 1400	ROA and LYH	No
10/02/80 0800	10/02/80 1330	LYH and DAN	No
06/03/82 0900	06/03/82 1530	LYH and DAN	No
09/15/82 1000	09/15/82 1300	LYH and DAN	No
10/06/82 1000	10/06/82 1500	LYH and DAN	No
11/02/82 1000	11/02/82 1530	LYH and DAN	No
11/03/82 0900	11/03/82 1830	ROA, LYH and DAN	No
11/23/82 0330	11/23/82 1600	ROA, LYH and DAN	No
04/08/83 1100	04/08/83 1500	ROA and LYH	No
06/30/83 0400	06/30/83 1300	ROA and LYH	No
07/01/83 0400	07/01/83 1300	ROA and LYH	No
11/02/83 1100	11/02/83 1500	ROA and DAN	No
11/03/95 0600	11/03/95 1300	ROA and LYH	No
11/07/96 0800	11/07/96 1200	LYH and DAN	No
11/27/01 0600	11/27/01 1600	ROA, LYH and DAN	No
09/03/02 0800	09/03/02 1330	LYH and DAN	No
10/23/02 0200	10/23/02 1600	LYH and DAN	No
11/21/02 2300	11/22/02 1400	LYH and DAN	No
08/21/03 1000	08/21/03 1300	ROA and LYH	No
05/18/04 0300	05/18/04 1230	LYH and DAN	No
09/12/04 1000	09/12/04 1400	LYH and DAN	No
06/30/05 0600	06/30/05 1100	LYH and DAN	No
01/13/06 0730	01/13/06 1930	LYH and DAN	No
07/26/06 1000	07/26/06 1400	LYH and DAN	No
11/14/08 0200	11/14/08 1230	ROA and LYH	No

#### IV. Radiation Post Cold Front Fog (34 cases)

Beginning	Ending	Locations with Fog	Precipitation
08/04/73 0800	08/04/73 1230	LYH and DAN	No
02/03/74 0800	02/03/74 1500	ROA, LYH and DAN	Yes
09/18/74 0100	09/18/74 1300	LYH and DAN	No
02/16/75 2200	02/17/75 1500	ROA, LYH and DAN	Yes
08/20/75 1000	08/20/75 1300	LYH and DAN	No
11/28/76 0800	11/28/76 2000	ROA, LYH and DAN	No
06/15/77 0700	06/15/77 1230	ROA and LYH	No
07/04/78 1000	07/04/78 1500	LYH and DAN	No
02/21/79 2300	02/22/79 1500	LYH and DAN	No
05/15/79 0800	05/15/79 1400	ROA, LYH and DAN	No
11/24/80 2300	11/25/80 0300	LYH and DAN	No
01/22/81 0000	01/22/81 1230	LYH and DAN	No
11/11/81 0700	11/11/81 1330	ROA and LYH	No
10/12/82 1030	10/12/82 1500	ROA and LYH	No
01/24/83 0000	01/24/83 0500	LYH and DAN	No
12/29/83 0000	12/29/83 0600	ROA and LYH	No
10/26/84 0300	10/26/84 1200	ROA, LYH and DAN	No
11/07/89 0200	11/07/89 1500	ROA, LYH and DAN	No
12/01/97 0100	12/01/97 1000	ROA, LYH and DAN	No
09/21/01 0400	09/21/01 0900	ROA and LYH	No
12/01/01 0700	12/01/01 0900	ROA, LYH and DAN	No
03/14/02 0230	03/14/02 1230	LYH and DAN	No
12/12/02 0200	12/12/02 1700	LYH and DAN	No
01/02/03 1200	01/02/03 2200	LYH and DAN	No
02/15/05 0100	02/15/05 1400	ROA, LYH and DAN	No
02/22/05 0600	02/22/05 1330	LYH and DAN	No
03/24/05 0330	03/24/05 0800	LYH and DAN	No
01/12/06 0000	01/12/06 1600	ROA, LYH and DAN	No
01/30/06 0700	01/30/06 1500	LYH and DAN	No
04/23/06 0200	04/23/06 1200	LYH and DAN	No
09/09/06 0630	09/09/06 1400	LYH and DAN	No
12/11/07 1130	12/11/07 1500	ROA and LYH	No
08/14/08 0300	08/14/08 1130	LYH and DAN	No
09/28/08 0500	09/28/08 1200	LYH and DAN	No

**V. Radiation Post Warm Front Fog (39 cases)**

Beginning	Ending	Locations with Fog	Measurable Precipitation
10/03/73 0900	10/03/73 1400	LYH and DAN	No
11/21/73 2330	11/22/73 1400	ROA, LYH and DAN	No
02/07/74 0500	02/07/74 1400	ROA and LYH	No
03/07/74 0700	03/07/74 1230	ROA and LYH	No
11/20/74 0830	11/20/74 1300	ROA and LYH	No
12/31/74 2330	01/01/75 0700	ROA and LYH	No
02/05/75 2300	02/06/75 1400	ROA and LYH	No
02/18/75 0230	02/18/75 1500	ROA, LYH and DAN	No
03/11/75 0000	03/11/75 1400	LYH and DAN	No
04/29/75 1000	04/29/75 1200	ROA, LYH and DAN	No
10/09/75 0500	10/09/75 1330	ROA and LYH	No
11/27/76 0800	11/27/76 1500	ROA and LYH	No
11/08/77 1100	11/08/77 1600	LYH and DAN	No
11/24/77 1100	11/24/77 1600	ROA, LYH and DAN	No
12/01/77 1100	12/01/77 1600	LYH and DAN	No
01/07/78 0000	01/07/78 1530	ROA and DAN	No
12/20/78 1600	12/21/78 0300	LYH and DAN	No
10/11/80 0530	10/11/80 1200	ROA and LYH	No
01/19/82 2200	01/20/82 1000	ROA, LYH and DAN	No
02/16/82 0530	02/16/82 1430	ROA and LYH	No
11/29/82 1000	11/29/82 1400	ROA and LYH	No
12/02/82 0000	12/02/82 1730	ROA, LYH and DAN	No
01/11/83 0030	01/11/83 0500	LYH and DAN	No
03/07/83 0130	03/07/83 1600	ROA, LYH and DAN	No
04/09/83 0900	04/09/83 1630	ROA and LYH	No
10/19/84 0900	10/19/84 1300	ROA, LYH and DAN	No
02/08/86 0400	02/08/86 1400	LYH and DAN	No
02/20/86 0200	02/20/86 1500	ROA, LYH and DAN	No
03/02/97 0000	03/02/97 0900	ROA, LYH and DAN	No
11/23/99 0300	11/23/99 1400	ROA and DAN	No
11/08/00 0600	11/08/00 1300	LYH and DAN	No
11/22/02 0030	11/22/02 1400	LYH and DAN	No
02/23/03 0000	02/23/03 0600	ROA and LYH	No
10/30/04 0400	10/30/04 1200	ROA and LYH	No
12/07/04 1100	12/07/04 1600	ROA, LYH and DAN	No
12/10/04 1400	12/10/04 1800	LYH and DAN	No
10/18/06 0630	10/18/06 1400	ROA, LYH and DAN	No
12/23/06 0600	12/23/06 1330	ROA and LYH	No
12/26/06 0500	12/26/06 1130	ROA and DAN	No

(CONTINUED FROM FRONT COVER)

- NWS ER 46 An Objective Method of Forecasting Summertime Thunderstorms. John F. Townsend and Russell J. Younkin. May 1972. (COM-72-10765).
- NWS ER 47 An Objective Method of Preparing Cloud Cover Forecasts. James R. Sims. August 1972. (COM-72-11382).
- NWS ER 48 Accuracy of Automated Temperature Forecasts for Philadelphia as Related to Sky Condition and Wind Direction. Robert B. Wassall. September 1972. (COM-72-11473).
- NWS ER 49 A Procedure for Improving National Meteorological Center Objective Precipitation Forecasts. Joseph A. Ronco, Jr. November 1972. (COM-73-10132).
- NWS ER 50 PEATMOS Probability of Precipitation Forecasts as an Aid in Predicting Precipitation Amounts. Stanley E. Wasserman. December 1972. (COM-73-10243).
- NWS ER 51 Frequency and Intensity of Freezing Rain/Drizzle in Ohio. Marvin E. Miller. February 1973. (COM-73-10570).
- NWS ER 52 Forecast and Warning Utilization of Radar Remote Facsimile Data. Robert E. Hamilton. July 1973. (COM-73-11275).
- NWS ER 53 Summary of 1969 and 1970 Public Severe Thunderstorm and Tornado Watches Within the National Weather Service, Eastern Region. Marvin E. Miller and Lewis H. Ramey. October 1973. (COM-74-10160).
- NWS ER 54 A Procedure for Improving National Meteorological Center Objective Precipitation Forecasts - Winter Season. Joseph A. Ronco, Jr. November 1973. (COM-74-10200).
- NWS ER 55 Cause and Prediction of Beach Erosion. Stanley E. Wasserman and David B. Gilhousen. December 1973. (COM-74-10036).
- NWS ER 56 Biometeorological Factors Affecting the Development and Spread of Plant Diseases. V.J. Valli. July 1974. (COM-74-11625/AS).
- NWS ER 57 Heavy Fall and Winter Rain In The Carolina Mountains. David B. Gilhousen. October 1974. (COM-74-11761/AS).
- NWS ER 58 An Analysis of Forecasters' Propensities In Maximum/Minimum Temperature Forecasts. I. Randy Racer. November 1974. (COM-75-10063/AS).
- NWS ER 59 Digital Radar Data and its Application in Flash Flood Potential. David D. Sisk. March 1975. (COM-75-10582/AS).
- NWS ER 60 Use of Radar Information in Determining Flash Flood Potential. Stanley E. Wasserman. December 1975. (PB250071/AS).
- NWS ER 61 Improving Short-Range Precipitation Guidance During the Summer Months. David B. Gilhousen. March 1976. (PB256427).
- NWS ER 62 Locally Heavy Snow Downwind from Cooling Towers. Reese E. Otts. December 1976. (PB263390/AS).
- NWS ER 63 Snow in West Virginia. Marvin E. Miller. January 1977. (PB265419/AS).
- NWS ER 64 Wind Forecasting for the Monongahela National Forest. Donald E. Risher. August 1977. (PB272138/AS).
- NWS ER 65 A Procedure for Spraying Spruce Budworms in Maine during Stable Wind Conditions. Monte Glovinsky. May 1980. (PB80-203243).
- NWS ER 66 Contributing Factors to the 1980-81 Water Supply Drought, Northeast U.S. Solomon G. Summer. June 1981. (PB82-172974).
- NWS ER 67 A Computer Calculation and Display System for SLOSH Hurricane Surge Model Data. John F. Townsend. May 1984. (PB84-198753).
- NWS ER 68 A Comparison Among Various Thermodynamic Parameters for the Prediction of Convective Activity. Hugh M. Stone. April 1985. (PB85-206217/AS).
- NWS ER 69 A Comparison Among Various Thermodynamic Parameters for the Prediction of Convective Activity, Part II. Hugh M. Stone. December 1985. (PB86-142353/AS).
- NWS ER 70 Hurricane Gloria's Potential Storm Surge. Anthony G. Gigi and David A. Wert. July 1986. (PB86-226644/AS).
- NWS ER 71 Washington Metropolitan Wind Study 1981-1986. Clarence Burke, Jr. and Carl C. Ewald. February 1987. (PB87-151908/AS).
- NWS ER 72 Mesoscale Forecasting Topics. Hugh M. Stone. March 1987. (PB87-180246/AS).
- NWS ER 73 A Procedure for Improving First Period Model Output Statistics Precipitation Forecasts. Antonio J. Lacroix and Joseph A. Ronco, Jr. April 1987. (PB87-180238/AS).
- NWS ER 74 The Climatology of Lake Erie's South Shoreline. John Kwiatkowski. June 1987. (PB87-205514/AS).
- NWS ER 75 Wind Shear as a Predictor of Severe Weather for the Eastern United States. Hugh M. Stone. January 1988. (PB88-157144).
- NWS ER 76 Is There A Temperature Relationship Between Autumn and the Following Winter? Anthony Gigi. February 1988. (PB88-173224).
- NWS ER 77 River Stage Data for South Carolina. Clara Cillentine. April 1988. (PB88-201991/AS).
- NWS ER 78 National Weather Service Philadelphia Forecast Office 1987 NOAA Weather Radio Survey & Questionnaire. Robert P. Wanton. October 1988. (PB89-111785/AS).
- NWS ER 79 An Examination of NGM Low Level Temperature. Joseph A. Ronco, Jr. November 1988. (PB89-122543/AS).
- NWS ER 80 Relationship of Wind Shear, Buoyancy, and Radar Tops to Severe Weather 1988. Hugh M. Stone. November 1988. (PB89-1222419/AS).
- NWS ER 81 Relation of Wind Field and Buoyancy to Rainfall Inferred from Radar. Hugh M. Stone. April 1989. (PB89-208326/AS).
- NWS ER 82 Second National Winter Weather Workshop, 26-30 Sept. 1988: Postprints. Laurence G. Lee. June 1989. (PB90-147414/AS).
- NWS ER 83 A Historical Account of Tropical Cyclones that Have Impacted North Carolina Since 1586. James D. Stevenson. July 1990. (PB90-259201).
- NWS ER 84 A Seasonal Analysis of the Performance of the Probability of Precipitation Type Guidance System. George J. Maglaras and Barry S. Goldsmith. September 1990. (PB93-160802).
- NWS ER 85 The Use of ADAP to Examine Warm and Quasi-Stationary Frontal Events in the Northeastern United States. David R. Vallee. July 1991. (PB91-225037).
- NWS ER 86 Rhode Island Hurricanes and Tropical Storms A Fifty-Six Year Summary 1936-1991. David R. Vallee. March 1993. (PB93-162006).
- NWS ER 87 Post-print Volume, Third National Heavy Precipitation Workshop, 16-20 Nov. 1992. April 1993. (PB93-186625).
- NWS ER 88 A Synoptic and Mesoscale Examination of the Northern New England Winter Storm of 29-30 January 1990. Robert A. Marine and Steven J. Capriola. July 1994. (PB94-209426).
- NWS ER 89 An Initial Comparison of Manual and Automated Surface Observing System Observations at the Atlantic City, New Jersey, International Airport. James C. Hayes and Stephan C. Kuhl. January 1995.
- NWS ER 90 Numerical Simulation Studies of the Mesoscale Environment Conducive to the Raleigh Tornado. Michael L. Kaplan, Robert A. Rozumalski, Ronald P. Weglarz, Yuh-Lang Lin, Steven Businger, and Rodney F. Gonski. November 1995.
- NWS ER 91 A Climatology of Non-convective High Wind Events in Western New York State. Thomas A. Niziol and Thomas J. Paone. April 2000.
- NWS ER 92 Tropical Cyclones Affecting North Carolina Since 1586 - An Historical Perspective. James E. Hudgins. April 2000.
- NWS ER 93 A Severe Weather Climatology for the Wilmington, NC WFO County Warning Area. Carl R., Morgan. October 2001.
- NWS ER 94 Surface-based Rain, Wind, and Pressure Fields in Tropical Cyclones over North Carolina since 1989. Joel Cline. June 2002.
- NWS ER 95 A Severe Weather Climatology for the Charleston, South Carolina, WFO County Warning Area. Stephen Brueske, Lauren Plourd, Matthen Volkmer. July 2002.
- NWS ER 96 A Severe Weather Climatology for the WFO Wakefield, VA County Warning Area. Brian T. Cullen. May 2003. (PB2003-105462).
- NWS ER 97 Severe Weather Climatology for the Columbia, SC WFO County Warning Area. Leonard C. Vaughan. September 2003. (PB2004-100999).
- NWS ER 98 Climatology of Heavy Rainfall Associated with Tropical Cyclones Affecting the Central Appalachians. James Hudgins, Steve Keighton, Kenneth Kostura, Jan Jackson. September 2005.

NWS ER 99 A Severe Weather Climatology for the WFO Blacksburg, Virginia, County Warning Area. Robert Stonefield, James Hudgins. January 2007.

NWS ER 100 Tropical Cyclones Affecting North Carolina Since 1586 - An Historical Perspective. James E. Hudgins. October 2007.

NWS ER 101 A Severe Weather Climatology for the Raleigh, NC County Warning Area. Clyde Brandon Locklear. May 2008.

NWS ER 102 A Severe Weather Climatology for the Wilmington, OH County Warning Area (1950-2004). Michael D. Ryan. May 2008.

NWS ER 103 A Climatology of Flash Flood Events for the National Weather Service Eastern Region, Alan M. Cope. June 2009.

NWS ER 104 An Abbreviated Flash Flood/Flood Climatology (1994-2007) for the WFO Blacksburg, VA County Warning Area., Robert Stonefield and Jan Jackson. September 2009.

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS**--Important definitive research results, major techniques, and special investigations.

**CONTRACT AND GRANT REPORTS**--Reports prepared by contractors or grantees under NOAA sponsorship.

**ATLAS**--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

**TECHNICAL SERVICE PUBLICATIONS**--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

**TECHNICAL REPORTS**--Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS**--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE  
U.S. DEPARTMENT OF COMMERCE  
5285 PORT ROYAL ROAD  
SPRINGFIELD, VA 22161